

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS ✓
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.


**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



Figs. 1A-1F

Fig. 2





Figs. 3A - 3B

Ac-R18L-NH₂ i.v. (100 μ g/mouse)
Apo E null, fasted




Fig. 4

Ac-R18L-NH₂, i.v. dose dependency
Apo E null; fasted





Fig. 5

24 h fecal total cholesterol
(collected from time of injection;
apo E null; 100 μ g R18L i.v.)



■ Control
■ R18L

Fig. 6






Fig. 7

REVIEWED BY
S. B. COOPER
15 JUN 1968
FBI - NEW YORK



R14-L2
Fig. 8B

R14 L-1
Fig. 8A



Uptake of LDL: Effect of R14L peptides




Fig. 9

**Apo E null; i.v. injections of two
single domain cationic peptides**




Fig. 10

Apo E null; 100 μ g peptide i.v.
Ac-(R)14L-NH₂ analog 2






Fig. 11

Figs. 12A-12F


Phase	Molécular structure
-------	---------------------




Bilayer



Cylindric



Micellar



Hexagonal (H_{11})



Cone